

# Transfection

+ How do you study  
and control gene expression?

+ How do you detect  
mycoplasma contamination?

+ How do you ensure the success  
of your RNA or protein transfection?

# Introduction to Transfection

Transfection is the delivery of DNA, RNA, proteins, and macromolecules into eukaryotic cells. Goals for transfection include the study of gene regulation as well as protein expression and function. The success of transfection depends on transfection efficiency, low cytotoxicity, and reproducibility.

To ensure high efficiency of your transfection, you need reliable transfection technology and reagents to optimize your cell culture conditions. We provide you with a variety of reagents for your preferred transfection method and *mycoplasma* detection kits to ensure early detection of cell culture contamination.

For more information on our transfection products, please visit  
[www.stratagene.com/transfection](http://www.stratagene.com/transfection).



## CONTENTS:

### Transfection

- I Transfection Selection Guide
- 2-3 Transient and Stable Transfections
- 4-5 Special Transfection Needs

### Mycoplasma

- 6 Mycoplasma Selection Guide
  - 7-8 Mycoplasma Detection
- 
- 9 Ordering Information

**Table 1: Selection Guide for Transfection Reagents**

|                                                    | RECOMMENDED APPLICATION                  | TRANSFECTION METHOD        | FEATURES                                                                                                                                                                         | INCUBATION TIME |
|----------------------------------------------------|------------------------------------------|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------|
| Mammalian Transfection Kit                         | DNA transfection                         | CaPO <sub>4</sub> , DEAE   | + Works for many cell types                                                                                                                                                      | + 12 - 24 hrs   |
| MBS Mammalian Transfection Kit                     | DNA transfection                         | Modified CaPO <sub>4</sub> | + Works with low DNA concentration<br>+ Recommended for CHO and HEK293 cells                                                                                                     | + 3 hrs         |
| LipoTaxi <sup>®</sup> Transfection Reagent         | DNA transfection                         | Liposome based             | + Works for many cell types<br>+ Requires small amounts of DNA<br>+ Gentle on cells<br>+ Room temperature storage                                                                | + 4 - 6 hrs     |
| GeneJammer <sup>®</sup> Transfection Reagent       | DNA transfection                         | Polyamine based            | + Good for hard to transfect (e.g. primary cells)<br>+ Works for many cell types<br>+ Effective in presence of serum (no media changes required)<br>+ Less cytotoxic than lipids | + 3 hrs         |
| GeneEraser <sup>™</sup> siRNA Transfection Reagent | siRNA transfection                       | Proprietary formulation    | + Optimized for siRNA transfection into many cell types<br>+ Knocks down target gene even at low siRNA concentration<br>+ Low cell cytotoxicity<br>+ Works in presence of serum  | + 20 - 72 hrs   |
| BioTrek <sup>™</sup> Protein Delivery Reagent      | Protein, peptide, macromolecule delivery | Lipid formulation          | + Works for many cell types<br>+ Conveniently packaged in single use vials<br>+ Includes control proteins                                                                        | + 4 hrs         |
| ViraPack <sup>™</sup> Transfection Kit             | Viral packaging                          | CaPO <sub>4</sub>          | + Optimized for viral packaging cell lines                                                                                                                                       | + 3 hrs         |




# Transient and Stable Transfections

Classical transfection techniques include the DEAE-dextran method, calcium phosphate method, and liposome-mediated transfection (lipofection). Your choice of transfection technology will influence your results. We offer a variety of transfection reagents and kits to suit your specific needs.


## DEAE-dextran and Calcium Phosphate Methods

We offer several mammalian transfection kits based on the original methods used to introduce DNA into cultured mammalian cells: DEAE-dextran and calcium phosphate. Our original Mammalian Transfection Kit utilizes the modified calcium phosphate method, which yields up to 100-fold efficiency over traditional methods, depending on the cell line to be transfected (Figure 1). Both are relatively simple techniques, with DEAE-dextran better suited for transient transfections and the calcium phosphate method for stable transfections.

To increase transfection efficiency over these standard methods, we utilize a specially modified bovine serum in our MBS Mammalian Transfection Kit<sup>1</sup> for use during transfection. This results in significantly higher efficiencies as well as reduced assay time (Figure 2). This kit exhibits superior performance with CHO (Chinese Hamster Ovary) and HEK293 (Human Embryonic Kidney) cells.



**Figure 1**  
INCREASED YIELD WITH MAMMALIAN TRANSFECTION KIT




**Figure 2**  
INCREASED EFFICIENCY WITH MBS MAMMALIAN TRANSFECTION KIT

### Lipofection

Compared to the DEAE-dextran and calcium phosphate methods, liposomes often offer higher transfection efficiency and better reproducibility. Using our LipoTAXI® Transfection Reagent results in stable and transient transfections with low cytotoxicity and consistent high transfection efficiencies (a 10-fold increase over calcium phosphate/DEAE mediated transfection in many cell lines). Furthermore, our LipoTAXI reagent can be conveniently stored at room temperature.


### Polyamines

Our GeneJammer® Transfection Reagent provides the high transfection efficiencies that you need, even in the most problematic conditions such as in the presence of serum or in transfection of primary cells (Figure 3). Furthermore, this easy-to-use reagent results in significantly reduced levels of cell damage as compared to the traditional liposomal method (Figure 4).



**Figure 3**  
**GENEJAMMER® REAGENT ENSURES HIGH TRANSFECTION EFFICIENCY IN BOTH EASY- AND DIFFICULT-TO-TRANSFECT CELLS**

We used pCMV-beta-galactosidase to transfect COS-7 cells (A) and human umbilical vein endothelial cells (B), using GeneJammer® reagent and competitor transfection reagents.



**Figure 4**  
**GENEJAMMER® REAGENT DEMONSTRATES LOWEST TOXICITY, HIGHEST NUMBER OF VIABLE CELLS**


To compare viability, we transfected HeLa cells with pCMV-beta-galactosidase and GeneJammer® reagent or competitive reagents. 24 hours after transfection, cell viability was evaluated by counting cells stained with trypan blue.

# Special Transfection Needs

We also offer a variety of specialty transfection reagents and kits to meet your experimental needs such as transfecting RNA, proteins, or for virus packaging.

## Protein Delivery

Our BioTrek™ Protein Delivery Reagent<sup>2</sup> allows you to directly translocate your protein into a living cell. Conveniently packaged in single use vials, this provides an easy method to effectively deliver proteins, peptides, and other macromolecules directly into a broad range of cell types at efficiencies of 50–95% in as little as 3–4 hours (Figure 5). This enables you to rapidly and accurately study cellular and protein function.



**Figure 5**

### PROTEIN TRANSFECTION OF BETA-GALACTOSIDASE WITH BIOTREK™ REAGENT

BioTrek™ reagent mediated intracellular delivery of beta-galactosidase in NIH-3T3 cells (A). After 4 hours, cells were fixed, stained for activity with the X-gal reagent, then compared to standard transfection (B).


siRNA Transfection

RNA interference (RNAi) is a powerful method for inhibiting the expression of a particular gene. RNAi experiments require a transfection reagent that is specially optimized for siRNA delivery into mammalian cells.

Our GeneEraser™ siRNA Transfection Reagent<sup>3</sup> enables the use of very low levels of siRNA to knock out stable or transient gene expression in a wide variety of cell lines and has a high transfection efficiency (Figure 6). Provided in a ready-to-use format, the GeneEraser reagent and your siRNA can be added directly to cells in media, even in the presence of serum.

High Titer Virus Production

The transfection efficiency of the packaging cell line in any viral delivery experiment is critical. Our ViraPack™ Transfection Kit<sup>1</sup> transfects the HEK293 packaging cell line with exceptional efficiency, so you can increase the number of infectious virus particles produced per transfection.



**Figure 6**  
**SILENCING EFFECT OF LUCIFERASE siRNA BY GENEERASER™ siRNA TRANSFECTION REAGENT**  
Hela cells were transfected with 10 µg luciferase reporter plasmid pGL2 (Promega) using our GeneJammer™ siRNA Transfection Reagent. Cells were then transfected with luciferase GL2 siRNA duplex (Dharmacon Research) using different transfection reagents (2.5 µl). 83 to 90% knock out of luciferase expression was observed when 2 to 30 nM concentration of siRNA was complexed with the GeneEraser reagent, compared to 40 to 60% knock out with Reagent A and 13 to 22% knock out with Reagent B.

**Table 2**  
The GeneEraser™ siRNA Transfection Reagent has been used to successfully deliver siRNA into the cell types listed above.

# Mycoplasma Detection

Present in 5 - 35% of cell cultures, *mycoplasma* contamination can alter growth characteristics, enzyme patterns, cell membrane composition, chromosomal structure, and transfection efficiency. These lead to unreliable experimental results as well as loss of time and precious cell lines. Therefore, routine screening for *mycoplasma* contamination is essential. Unlike bacteria and fungi, *mycoplasma* contamination cannot be detected by visual inspection. Their small size allows them to pass through most sterilization filters and they are resistant to common antibiotics.



Table 3: Mycoplasma Selection Guide

|                                 | METHOD         | FEATURES                                                                                                                                                                                                                                                                                                                                   | RECOMMENDED APPLICATION |
|---------------------------------|----------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------|
| Mycoplasma Plus™ PCR Primer Set | PCR            | + Includes positive and internal controls                                                                                                                                                                                                                                                                                                  | + Detection, speciation |
| MycoSensor™ PCR Assay Kit       | PCR            | + Primer design prevents detection of <i>E. coli</i> DNA that often contaminates recombinant <i>Taq</i> DNA polymerase preparations<br><br>+ Short cycle time (50% faster)<br><br>+ Protocol ensures decontamination of carry-over PCR product<br><br>+ Detects as few as 10 copies of <i>mycoplasma</i> genomic DNA, depending on species | + Detection             |
| MycoSensor™ QPCR Assay Kit      | Real-Time QPCR | + Quick results (2 hours)<br><br>+ Convenient master mix format<br><br>+ Detects as few as 10 copies of <i>mycoplasma</i> genomic DNA, depending on species<br><br>+ Includes positive and internal amplification controls<br><br>+ Includes PCR purification kit<br><br>+ Tested on multiple real-time QPCR platforms                     | + Detection             |



Mycoplasma Detection

Traditional *mycoplasma* detection methods, such as culture testing, staining, and immunofluorescence detection, are very time-consuming, not sensitive, and difficult to interpret. PCR amplification provides a quick and easy method for *mycoplasma* detection.

The Mycoplasma Plus™ PCR Primer Set provides you reliable PCR detection and speciation of *mycoplasma*. This kit provides an internal control to confirm PCR amplification was successful and a noninfectious *mycoplasma* genomic DNA template that serves as a positive control.

Our next generation PCR amplification kit, the MycoSensor™ PCR Assay Kit<sup>‡</sup>, features enhanced primer design to improve detection of the eight most common contaminating species (Table 4) and to prevent the detection of *E. coli* DNA, which often contaminates recombinant *Taq* DNA polymerase preparations. The MycoSensor kit features shorter run time (50% faster) and detects as few as 10 copies of *Mycoplasma* genomic DNA, depending on species (Figure 7) .




Figure 7  
ELIMINATING FALSE NEGATIVES IN YOUR MYCOPLASMA DETECTION

The internal control in the MycoSensor™ PCR Assay Kit reduces false negatives by verifying that PCR amplification was successful, indicated by a 500-bp band that is distinguishable from the *mycoplasma* target band.  
**Lane 1:** kb ladder; **Lane 2:** Water + internal control (no template); **Lane 3:** Internal control + *M. orale* (Positive control); **Lane 4:** Internal control + *A. laidlawii* (Positive control); **Lane 5:** Internal control + CHO cell supernatant (Mycoplasma -); **Lane 6:** Internal control + HeLa cell supernatant (Mycoplasma +).

EIGHT MOST COMMON SPECIES OF MYCOPLASMA AND ACHOLEPLASMA

|   |                      |
|---|----------------------|
| 1 | <i>A. laidlawii</i>  |
| 2 | <i>M. arginini</i>   |
| 3 | <i>M. fermentans</i> |
| 4 | <i>M. hominis</i>    |
| 5 | <i>M. hyorhina</i>   |
| 6 | <i>M. orale</i>      |
| 7 | <i>M. pirum</i>      |
| 8 | <i>M. salivarium</i> |

Table 4  
Our MycoSensor™ PCR Assay Kit detects the eight most common contaminant species of *mycoplasma* and *acholeplasma*.


### Mycoplasma Detection by Real-Time PCR

Our new MycoSensor™ QPCR Assay Kit<sup>5</sup> detects *mycoplasma* infection of cell cultures by real-time quantitative PCR, using SYBR® Green dye detection, in less than 2 hours. The convenient master mix format provides all the reagents needed for amplification and fluorescence detection, and the kit detects the eight most common cell culture contaminating species of *mycoplasma* (Table 4). Customers can detect as few as 10 copies of contaminating genomic DNA from 100 µl of cell culture supernatant, or test for weak *mycoplasma* infection in extracts made from cell pellets (Figure 8).

The kit includes two positive control templates to validate the detection of successful PCR amplification of *mycoplasma* template and confirm the SYBR® Green dissociation profile for your test samples. If your test cell line is infected, the SYBR Green dissociation profile results in a melting peak at 82°C. As seen by a melting peak around 85°C, the amplification control confirms the absence of PCR inhibitors, thus reducing false negative results (Figure 9). To minimize false positives,


the closed tube real-time detection format minimizes the potential for cross-contamination with PCR amplicon. We include a passive reference dye in the kit to allow you the flexibility of running the assay on most real-time QPCR platforms. The kit also contains DNA purification reagents for removal of potential PCR inhibitors in cell culture supernatants and cell pellets.

The MycoSensor QPCR assay kit provides a fast, convenient method for the routine testing of cell cultures for *mycoplasma* and *acholeplasma* contamination. The kit delivers consistent results and includes controls designed to eliminate false negatives and false positives. We've optimized the MycoSensor QPCR assay kit for maximum performance on multiple real-time QPCR platforms, including our Mx3000P™ and Mx4000® Quantitative PCR instruments, the Cepheid Smart Cycler®, and the ABI PRISM® 7700.



**Figure 8**  
DETECTION OF *M. ORALE* GENOMIC DNA USING MYCOSENSOR™ QPCR ASSAY KIT

The MycoSensor™ QPCR Assay Kit detects a range from 100,000 copies to 10 copies of *M. orale* genomic DNA in the presence of HeLa cell culture supernatant. The reactions were done in duplicate on the Mx3000P™ system and the amplification plots are shown here.



**Figure 9**  
SPECIFIC DETECTION OF MYCOPLASMA CONTAMINATION AND AMPLIFICATION CONTROL

The SYBR® Green dissociation curve generated on the Mx3000P™ system for the *mycoplasma* primers and the amplification control primers indicates distinct melting temperatures of ~82°C for mycoplasma and ~85°C for the amplification control, respectively.

# Ordering Information

| Description                                     | Catalog No.                   | Size   | Price (\$) |
|-------------------------------------------------|-------------------------------|--------|------------|
| <b>Mycoplasma Detection Kits</b>                |                               |        |            |
| MYCOPLASMA PLUS™ PCR PRIMER SET                 | 1 kit (100 rxn)               | 302008 | 335        |
| MYCOSENSOR™ PCR ASSAY KIT                       | 1 kit (100 rxn)               | 302108 | 335        |
| MYCOSENSOR™ QPCR ASSAY KIT                      | 1 kit (100 rxn)               | 302106 | 395        |
| <b>Transfection Reagents &amp; Kits</b>         |                               |        |            |
| GENEJAMMER® TRANSFECTION REAGENT                | 0.3 ml                        | 204132 | 102        |
|                                                 | 1.0 ml                        | 204130 | 202        |
|                                                 | 4 x 1.0 ml                    | 204131 | 728        |
| LIPOTAXI® TRANSFECTION REAGENT                  | 3 x 1 ml                      | 204110 | 211        |
| <b>MAMMALIAN TRANSFECTION KITS</b>              |                               |        |            |
| CAPO4 AND DEAE                                  | 1 kit (30 transfections each) | 200285 | 237        |
| CAPO4                                           | 1 kit (30 transfections)      | 200385 | 113        |
| DEAE                                            | 1 kit (30 transfections)      | 200386 | 113        |
| MBS MAMMALIAN TRANSFECTION KIT                  | 1 kit                         | 200388 | 330        |
| <b>Protein Delivery</b>                         |                               |        |            |
| BIOTREK™ PROTEIN DELIVERY REAGENT               | 1 kit (24 rxn)                | 204140 | 270        |
| <b>siRNA Transfection</b>                       |                               |        |            |
| GENEERASER™ SIRNA TRANSFECTION REAGENT          | 0.4 ml                        | 204152 | 177        |
|                                                 | 1.0 ml                        | 204150 | 358        |
| <b>Viral Delivery</b>                           |                               |        |            |
| VIRAPACK™ TRANSFECTION KIT                      | 1 kit                         | 200488 | 190        |
| <b>Other Related Products</b>                   |                               |        |            |
| BETA-GALACTOSIDASE ASSAY KIT                    | 1 kit (100 rxn)               | 200383 | 149        |
| BETA-GALACTOSIDASE ASSAY KIT (HIGH SENSITIVITY) | 1 kit (100 rxn)               | 200710 | 180        |
| BETA-GALACTOSIDASE ASSAY KIT (IN SITU)          | 1 kit                         | 200384 | 154        |
| G418 SULFATE                                    | 100 mg                        | 200397 | 27         |
|                                                 | 500 mg                        | 200398 | 52         |
|                                                 | 1 mg                          | 200399 | 93         |
|                                                 | 10 ml                         | 200049 | 62         |
| HYGROMYCIN B                                    | 1 MU (1 x 10 <sup>6</sup> U)  | 200715 | 129        |
| LUCIFERASE ASSAY KIT                            | 1 kit (100 rxn)               | 219020 | 90         |
| QUANTOS CELL PROLIFERATION ASSAY KIT            | 1 kit                         | 302011 | 221        |

- LEGAL INFORMATION
- 1

U.S. Patent No. 5,330,904
- 2

BioTrek™ Protein Delivery Reagent is manufactured for Stratagene by Gene Therapy Systems, Inc.
- 3

GeneEraser™ siRNA Transfection Reagent is manufactured for Stratagene by Mirus
- 4

Patent pending

Purchase of this PCR-related product does not convey any rights under the PCR patents owned by Hoffmann-La Roche. A license to use the PCR process accompanies the purchase of certain reagents from Stratagene when used in conjunction with an Authorized Thermal Cycler.
- 5

Patent pending

Purchase of these products is accompanied by a license to use them in the Polymerase Chain Reaction (PCR) process in conjunction with a thermal cycler whose use in the automated performance of the PCR process is covered by the up-front license fee, either by payment to Applied Biosystems or as purchased, i.e., an authorized thermal cycler.
- SYBR® is a registered trademark of Molecular Probes

FuGENE 6 is a registered trademark of Fugent, LLC, USA

GenePORTER 2 is a trademark of Gene Therapy Systems, Inc.

Lipofectamine, Lipofectamine Plus, and Lipofectamine 2000 are trademarks of Invitrogen Corp.

SuperFect is a registered trademark of Qiagen Inc.

SYBR® is a registered trademark of Molecular Probes
- www.stratagene.com

AMPLIFICATION

CELL BIOLOGY

CLONING

MICROARRAYS

NUCLEIC ACID  
ANALYSIS

PROTEIN FUNCTION  
& ANALYSIS

QUANTITATIVE  
PCR

SOFTWARE  
SOLUTIONS

### Stratagene USA and Canada

Order: 800-424-5444 x3

Technical Services: 800-894-1304 x2

### Stratagene Europe

Order: 00800-7000-7000

Technical Services: 00800-7400-7400

### Stratagene Japan K.K.

Order: 3-5821-8077

Technical Services: 3-5821-8076

### Distributors >

For a list of worldwide distributors, please visit:

[www.stratagene.com](http://www.stratagene.com)

© Stratagene, an Agilent Technologies company, 2008  
Printed in USA, April 2008  
BR61 5989-8234EN

# STRATAGENE

An **Agilent Technologies** Company

11011 North Torrey Pines Road  
La Jolla, CA 92037 USA

PRSRT STD  
U.S. POSTAGE  
**PAID**  
San Diego, CA  
Permit No. 1