

Comparison of Denaturing and Nondenaturing Mobile Phases in SEC: ZORBAX GF-250,


+/- GuCI

Application

Biochemical

Robert Ricker

Size-exclusion chromatography is a powerful technique for the size-dependent separation of biomolecules. While standard, non-denaturing mobile phases are often desirable, denaturing mobile phases containing guanidine hydrochloride (or SDS), may be used to insure un-aggregated, monomers for proper size determination. As shown below, denaturation shifts molecules toward larger apparent size and smaller retention volume.



Column: ZORBAX GF-250 (9.4 x 250), Agilent Part No. 884973-901

I.C: Hewlett Packard 1050

Flow: 2.0 mL / min.; ambient Inj. Vol.: 20 μ L (1 μ g / μ L)

Highlights

- Agilent ZORBAX GF-250 columns separate biomolecules in a sizedependent manner within a molecularweight range of 4,000-400,000 daltons. Note, some molecules (depending on their characteristics) are retained by non-SEC mechanisms and are shifted from linearity.
- ZORBAX GF-250 columns are manufactured using extremely hard particles, allowing high flow rates and fast separations.
- When larger molecules are used in combination with denaturing mobile phases, ZORBAX GF-450 may be used to obtain more-linear separation in the upper molecular-weight range.



Robert Ricker is an application chemist based at Agilent Technologies, Wilmington, Delaware.

For more information on our products and services, visit our website at: www.agilent.com/chem

Copyright[©] 2002 Agilent Technologies, Inc. All Rights Reserved. Reproduction, adaptation or translation without prior written permission is prohibited, except as allowed under the copyright laws.

Agilent shall not be liable for errors contained herein or for incidental or consequential damages in connection with the furnishing, performance, or use of this material.

Information, descriptions, and specifications in this publication are subject to change without notice.

Printed in the USA April 25, 2002 5988-6331EN

